

CY 2015 UDS Clinical Measures

Table, Line / Column	Clinical Measure	Brief Description	Universe (Denominator)	Performance Standard (Numerator)	Exclusions
6B, 7-9	Trimester of Entry into Prenatal Care	Prenatal care patients who entered prenatal care during their first trimester	Total number of women seen for prenatal care during the measurement year	Number of women entering prenatal care at the health center or with the referred provider during their first trimester	None
6B, 10	Childhood Immunization	Children who have received age appropriate vaccines prior to their 3rd birthday	Number of all children who: <ul style="list-style-type: none"> • were born between January 1, 2012, and December 31, 2012, • had at least one medical visit during the reporting period, • were first seen prior to their 3rd birthday 	Number of children fully immunized before their third birthday. "Fully immunized" includes: <ul style="list-style-type: none"> • Received ALL the following: 4 DTP/DTaP, 3 IPV, 1 MMR, 3 Hib, 3 HepB, 1 VZV (Varicella), and 4 Pneumococcal conjugate • OR evidence of contraindication for the vaccine or a history of having had the target illness 	None
6B, 11	Cervical Cancer Screening	Female patients aged 24-64 who received one or more Pap tests to screen for cervical cancer	Number of female patients: <ul style="list-style-type: none"> • born between January 1, 1951, and December 31, 1991 • who had at least one medical visit in a clinical setting 	Number of female patients 24-64 years of age receiving: <ul style="list-style-type: none"> • one or more Pap tests in a 3-year period from 2013 through 2015 • OR one or more Pap tests in a 5-year period from 2011 through 2015 IF they were 30 years of age or older at the time of the last Pap test <i>and</i> had a Pap test and an HPV test done simultaneously 	Had a hysterectomy and who have no residual cervix

CY 2015 UDS Clinical Measures

Table, Line / Column	Clinical Measure	Brief Description	Universe (Denominator)	Performance Standard (Numerator)	Exclusions
6B, 12	Weight Assessment and Counseling for Children and Adolescents	Children and adolescents aged 3 through 17 during measurement year with a BMI percentile, <i>and</i> counseling on nutrition and physical activity documented for the measurement year	Number of patients who: <ul style="list-style-type: none"> • were born between January 1, 1998, and December 31, 2012, • had at least one medical visit in a clinical setting, • and were seen ever by the health center prior to their 18th birthday 	Number of patients who: <ul style="list-style-type: none"> • had their BMI percentile (not just BMI or height and weight) documented during the measurement year, • AND had documentation of counseling for nutrition, • AND who had documentation of counseling for physical activity 	Pregnant patients
6B, 13	Adult Weight Screening and Follow-Up	Patients aged 18 and older with (1) BMI charted <i>and</i> (2) follow-up plan documented <i>if</i> patients are overweight or underweight	Number of patients who: <ul style="list-style-type: none"> • were born on or before December 31, 1997, • had at least one medical visit in a clinical setting, • and were last seen after their 18th birthday 	Number of patients who: <ul style="list-style-type: none"> • had their BMI (not just height and weight) documented during their last visit or within 6 months prior to that visit, • AND received a follow-up plan to address their weight if: <ul style="list-style-type: none"> ○ under age 65 and BMI ≥ to 25 or < 18.5 ○ age 65 or older and BMI ≥ 30 or < 23 	Pregnant women and terminally ill patients

CY 2015 UDS Clinical Measures

Table, Line / Column	Clinical Measure	Brief Description	Universe (Denominator)	Performance Standard (Numerator)	Exclusions
6B, 14a	Tobacco Use Screening and Cessation Intervention	Patients aged 18 and older who (1) were screened for tobacco use one or more times in the measurement year or the prior year <i>and</i> (2) for those found to be a tobacco user, received cessation counseling intervention or medication	Number of patients who: <ul style="list-style-type: none"> • were born on or before December 31, 1997, • had at least one medical visit during the reporting year, • had at least two medical visits ever, • and were seen after their 18th birthday 	Number of patients: <ul style="list-style-type: none"> • queried about their tobacco use one or more times during their most recent visit OR within 24 months of most recent visit, • AND, if found to be a tobacco user, received an order for smoking cessation medication or were on a cessation agent or received counseling 	None
6B, 16	Asthma Pharmacologic Therapy	Patients aged 5 through 40 diagnosed with persistent asthma who have an acceptable pharmacological treatment plan	Number of patients who: <ul style="list-style-type: none"> • had current diagnosis of persistent asthma • were born on or after January 1, 1975, and on or before December 31, 2010, and • had at least one medical visit during the reporting year, • had at least two visits ever, • and were last seen while age 5 through 40 	Number of patients who received a prescription for or were using: <ul style="list-style-type: none"> • inhaled corticosteroid, • OR an acceptable pharmacological agent: inhaled steroid combinations, anti-asthmatic combinations, antibody inhibitor, leukotriene modifiers, mast cell stabilizers, or methylxanthines 	Allergic reaction to asthma medications and those diagnosed with intermittent mild asthma

Table, Line / Column	Clinical Measure	Brief Description	Universe (Denominator)	Performance Standard (Numerator)	Exclusions
6B, 17	Coronary Artery Disease (CAD): Lipid Therapy	Patients aged 18 and older with a diagnosis of CAD who were prescribed a lipid-lowering therapy	<p>Number of patients who:</p> <ul style="list-style-type: none"> had an active diagnosis of CAD or diagnosed as having had a myocardial infarction (MI) or had cardiac surgery in the past were born on or before December 31, 1997, had at least one medical visit during the reporting year, had at least two medical visits ever, and were last seen after their 18th birthday 	Number of patients in the denominator who received a prescription for or were provided or were taking lipid lowering medications	Individuals whose last LDL lab test during the measurement year was less than 130 mg/dL and those with an allergy to or a history of adverse outcomes from or intolerance to LDL lowering medications
6B, 18	Ischemic Vascular Disease (IVD): Aspirin or Anti-Thrombotic Therapy	Patients aged 18 and older with a diagnosis of IVD or AMI, CABG, or PTCA procedure with aspirin or another antithrombotic therapy	<p>Number of patients who:</p> <ul style="list-style-type: none"> had an active diagnosis of IVD during the current or prior year OR had been discharged after AMI or CABG or PTCA in the prior year were born on or before December 31, 1997, had at least one medical visit during the reporting year, and were last seen after their 18th birthday 	Number of patients who had received a prescription for, were given, or were using Aspirin or another anti-thrombotic drug	None

CY 2015 UDS Clinical Measures

Table, Line / Column	Clinical Measure	Brief Description	Universe (Denominator)	Performance Standard (Numerator)	Exclusions
6B, 19	Colorectal Cancer Screening	Patients aged 51 through 74 years of age during measurement year with appropriate screening for colorectal cancer	Number of patients who were: <ul style="list-style-type: none"> • born between January 1, 1941, and December 31, 1964, • and had at least one medical visit during the reporting year 	Number of patients with appropriate screening for colorectal cancer: <ul style="list-style-type: none"> • a colonoscopy after January 1, 2006, • OR a flexible sigmoidoscopy after January 1, 2011, • OR a fecal occult blood test (FOBT), including the fecal immunochemical (FIT) test during the measurement year 	Had colorectal cancer or colectomy
6B, 20	HIV Linkage to Care	Patients newly diagnosed with HIV were seen for follow-up treatment within 90 days of diagnosis	Number of patients: <ul style="list-style-type: none"> • first diagnosed with HIV by the health center between October 1 of prior year through September 30 of the current measurement year • and had at least one medical visit during the measurement year or prior year 	Number of newly diagnosed HIV patients that received treatment with 90 days of diagnosis from a medical provider OR had received medical treatment from a referral resource	None

CY 2015 UDS Clinical Measures

Table, Line / Column	Clinical Measure	Brief Description	Universe (Denominator)	Performance Standard (Numerator)	Exclusions
6B, 21	Patients Screened for Depression and Follow-Up	Patients aged 12 and older who were screened for depression with a standardized tool and if screening was positive had a follow-up plan	Number of patients who: <ul style="list-style-type: none"> were born on or before December 31, 2003, and had at least one medical visit during the reporting year 	Number of patients who were screened for depression with a standardized tool and: <ul style="list-style-type: none"> test was negative, or screened positive for depression and had additional screening or a follow-up plan documented 	Patients with an active diagnosis for depression or bipolar disorder or already participating in on-going treatment for depression
6B, 22	Dental Sealants	Patients aged 6 through 9 with moderate to high risk for caries who received a sealant on a first permanent molar	Number of patient who: <ul style="list-style-type: none"> were at moderate to high risk for caries were born between January 1, 2006, and December 31, 2009, and had at least one oral assessment or periodic oral evaluation visit during the reporting year 	Number of patients who had a sealant on a first permanent molar during the measurement year	Children for whom all first permanent molars are non-sealable (i.e., molars are either decayed, filled, currently sealed, or un-erupted / missing)
7, 1b -1d	Low Birthweight	Babies, of health center prenatal care patients, born whose birthweight was below normal (less than 2500 grams).	Number of children born during measurement year to prenatal care patients	Number of each child born with a birthweight under 2500 grams	Sill-births or miscarriages

CY 2015 UDS Clinical Measures

Table, Line / Column	Clinical Measure	Brief Description	Universe (Denominator)	Performance Standard (Numerator)	Exclusions
7, 2a -2c	Controlled Hypertension	Patients diagnosed with hypertension (HTN) whose blood pressure (BP) was less than 140/90 at the time of the last reading	Number of patients who: <ul style="list-style-type: none"> • were first diagnosed as hypertensive before June 30 • were born between January 1, 1931, and December 31, 1997, • and have been seen for medical visits at least twice during the measurement year 	Number of patients whose last systolic blood pressure measurement was less than 140 mm Hg and whose last diastolic blood pressure was less than 90 mm Hg	Pregnant patients and patients with end state renal disease (ESRD)
7, 3a – 3d	Uncontrolled Diabetes	Patients with a diagnosis of Type I or Type II diabetes, whose hemoglobin A1c (HbA1c) was greater than 9%, or was missing a result, or an HbA1c test was not done during the measurement year	Number of patients who: <ul style="list-style-type: none"> • were diagnosed with diabetes • were born between January 1, 1941, and December 31, 1997, • and have been seen for medical visits at least twice during the measurement year 	Number of patients whose most recent hemoglobin A1c level during the measurement year is greater than 9% or did not have a test during the year or whose test result was missing	Patients with gestational diabetes, polycystic ovaries, or steroid-induced diabetes