



For Electronic Measure Specification Information go to: http://www.cms.gov/QualityMeasures/03_ElectronicSpecifications.asp#TopOfPage

| NQF Measure Number & PQRI Implementation Number | Clinical Quality Measure Title & Description | Clinical Quality Measure Steward & Contact Information | Core Clinical Quality Measure | UDS Overlap | Health Home Measure | NCQA Diabetes Recognition Program Measures |
|---|---|--|-------------------------------|---|--|--|
| NQF 0421 PQRI 128 | Title: Adult Weight Screening and Follow-Up Description: Percentage of patients aged 18 years and older with a calculated BMI in the past six months or during the current visit documented in the medical record AND if the most recent BMI is outside parameters, a follow-up plan is documented. | CMS/Quality Insights of Pennsylvania (QIP) Contact Information: www.usqualitymeasures.org | Core | 2011 UDS Measure. Table 6b, Section F, row 13. | NA | NA |
| NQF 0013 | Title: Hypertension: Blood Pressure Measurement Description: Percentage of patient visits for patients aged 18 years and older with a diagnosis of hypertension who have been seen for at least 2 office visits, with blood pressure (BP) recorded. | AMA-PCPI Contact Information: cpe@ama-assn.org | Core | 2011 UDS Measure. Similar to Table 7, Section B Hypertension is reported for all racial and ethnic groups. | NA | NA |
| NQF 0028 | Title: Preventive Care and Screening Measure Pair: a. Tobacco Use Assessment, b. Tobacco Cessation Intervention Description: Percentage of patients aged 18 years and older who have been seen for at least 2 office visits who were queried about tobacco use one or more times within 24 months b. Percentage of patients aged 18 years and older identified as tobacco users within the past 24 months and have been seen for at least 2 office visits, who received cessation intervention. | AMA-PCPI Contact Information: cpe@ama-assn.org | Core | 2011 UDS Measure. Table 6B, Section G, Row 14 and 15. | NA | NA |
| NQF 0024 | Title: Weight Assessment and Counseling for Children and Adolescents Description: Percentage of patients 2 -17 years of age who had an outpatient visit with a Primary Care Physician (PCP) or OB/GYN and who had evidence of BMI percentile documentation, counseling for nutrition and counseling for physical activity during the measurement year. | NCQA Contact Information: www.ncqa.org | Alternate Core | 2011 UDS Measure. Table 6b, Section E, row 12. | NA | NA |
| NQF 0038 | Title: Childhood Immunization Status Description: Percentage of children 2 years of age who had four diphtheria, tetanus and acellular pertussis (DTaP); three polio(IPV), one measles, mumps and rubella (MMR); two H influenza type B (HiB); three hepatitis B (Hep B); one chicken pox (VZV); four pneumococcal conjugate (PCV); two hepatitis A (Hep A); two, or three rotavirus (RV); and two influenza (flu) vaccines by their second birthday. The measure calculates a rate for each vaccine and nine separate combination rates | NCQA Contact Information: www.ncqa.org | Alternate Core | 2011 UDS Measure. Similar to Table 6b Section C, row 10. Childhood Immunization. Need to document each immunization discretely in order to capture on MU Reports. | NA | NA |
| NQF 0041 PQRI 110 | Title: Preventive Care and Screening: Influenza Immunization for Patients ≥ 50 Years Old Description: Percentage of patients aged 50 years and older who received an influenza immunization during the flu season (September through February). | AMA-PCPI Contact Information: cpe@ama-assn.org | Alternate Core | 2010 UDS Measure. Removed in 2011. | NA | NA |
| NQF 0059 PQRI 1 | Title: Diabetes: Hemoglobin A1c Poor Control Description: Percentage of patients 18 - 75 years of age with diabetes (type 1 or type 2) who had hemoglobin A1c >9.0% | National Committee for Quality Assurance (NCQA) Contact Information: www.ncqa.org | NA | 2011 UDS Measure. Similar to Table 7, Section C, Diabetes. | Not an Exact Match to NQF 0059 Hedis 2012 Effectiveness of Care Comprehensive Diabetes Care (HbA1c test and LDL-test) Percentage of members with diabetes who had at least one HbAc test and at least one LDL-C test | Title: HbA1c Testing & Control Proportion of patient panel with HbA1c >9.0%. Description: The proportion of patients 18 - 75 years of age with diabetes (type 1 or type 2) who had HbA1c >9.0%. Date and value of most recent HbA1c performed within the 12-month abstraction period. |
| NQF 0064 PQRI 2 | Title: Diabetes: Low Density Lipoprotein (LDL) Management and Control Description: Percentage of patients 18-75 years of age with diabetes (type 1 or type 2) who had LDL-C < 100 mg/dL. | NCQA Contact Information: www.ncqa.org | NA | NA | Not an exact match to NQF 0064 Hedis 2012 Effectiveness of Care Comprehensive Diabetes Care (HbA1c test and LDL-test) Percentage of members with diabetes who had at least one HbAc test and at least one LDL-C test | Title: Lipid Control Description: Proportion of patients 18-75 years of age with diabetes (type 1 or type 2) who had LDL ≥130mg/dL or < 100 mg/dL. Date and value of most recent LDL performed within the 12-month abstraction period. |

Note: Each provider must report on 3 core measures (or alternate core) and 3 additional measures.

| NQF Measure Number & PQRI Implementation Number | Clinical Quality Measure Title & Description | Clinical Quality Measure Steward & Contact Information | Core Clinical Quality Measure | UDS Overlap | Health Home Measure | NCQA Diabetes Recognition Program Measures |
|---|--|---|-------------------------------|--|---|--|
| NQF 0061 PQRI 3 | Title: Diabetes: Blood Pressure Management Description: Percentage of patients 18 - 75 years of age with diabetes (type 1 or type 2) who had blood pressure <140/90 mmHg. | NCQA Contact Information: www.ncqa.org | NA | NA | NA | Title: Blood Pressure Measurement Description: Proportion of patients 18-75 years of age with diabetes (type 1 or type 2) who had BP >140/90 mm HG or <130/80 mm Hg. Date and value of most recent BP measurement performed within the 12-month abstraction period. |
| NQF 0081 PQRI 5 | Title: Heart Failure (HF): Angiotensin-Converting Enzyme (ACE) Inhibitor or Angiotensin Receptor Blocker (ARB) Therapy for Left Ventricular Systolic Dysfunction (LVSD) Description: Percentage of patients aged 18 years and older with a diagnosis of heart failure and LVSD (LVEF < 40%) who were prescribed ACE inhibitor or ARB therapy. | American Medical Association-sponsored Physician Consortium for Performance Improvement (AMA-PCPI) Contact Information: cpe@ama-assn.org | NA | NA | Annual Monitoring for Patients on Persistent Medications (ACE/ARB and Diuretics) Hedis 2012- Effectiveness of Care: % of members who had 180 days of treatment with ambulatory medication therapy for ACE/ARB or Diuretics and who had the appropriate therapeutic monitoring | NA |
| NQF 0070 PQRI 7 | Title: Coronary Artery Disease (CAD): Beta-Blocker Therapy for CAD Patients with Prior Myocardial Infarction (MI) Description: Percentage of patients aged 18 years and older with a diagnosis of CAD and prior MI who were prescribed beta-blocker therapy. | AMA-PCPI Contact Information: cpe@ama-assn.org | NA | NA | Persistence of Beta Blocker Treatment after Heart Attack: Hedis 2012-Effectiveness of Care: % of members who were hospitalized and discharged alive with a diagnosis of AMI and who received persistent beta-blocker treatment for six months after discharge | NA |
| NQF 0043 PQRI 111 | Title: Pneumonia Vaccination Status for Older Adults Description: Percentage of patients 65 years of age and older who have ever received a pneumococcal vaccine. | NCQA Contact Information: www.ncqa.org | NA | NA | NA | NA |
| NQF 0031 PQRI 112 | Title: Breast Cancer Screening Description: Percentage of women 40-69 years of age who had a mammogram to screen for breast cancer. | NCQA Contact Information: www.ncqa.org | NA | 2011 UDS Measure on Table 6A - # of mammograms | NA | NA |
| NQF 0034 PQRI 113 | Title: Colorectal Cancer Screening Description: Percentage of adults 50-75 years of age who had appropriate screening for colorectal cancer. | NCQA Contact Information: www.ncqa.org | NA | 2012: New measure in Table 6B, Section K, Line 19. | Not an exact match to NQF 0034 Hedis 2012-Effectiveness of Care: % of member 50 and older who had appropriate screening for colorectal cancer | NA |
| NQF 0067 PQRI 6 | Title: Coronary Artery Disease (CAD): Oral Antiplatelet Therapy Prescribed for Patients with CAD Description: Percentage of patients aged 18 years and older with a diagnosis of CAD who were prescribed oral antiplatelet therapy. | AMA-PCPI Contact Information: cpe@ama-assn.org | NA | NA | NA | NA |
| NQF 0083 PQRI 8 | Title: Heart Failure (HF): Beta-Blocker Therapy for Left Ventricular Systolic Dysfunction (LVSD) Description: Percentage of patients aged 18 years and older with a diagnosis of heart failure who also have LVSD (LVEF < 40%) and who were prescribed beta-blocker therapy. | AMA-PCPI Contact Information: cpe@ama-assn.org | NA | NA | NA | NA |
| NQF 0105 PQRI 9 | Title: Anti-depressant medication management: (a) Effective Acute Phase treatment, (b)Effective Continuation Phase Treatment Description: The percentage of patients 18 years of age and older who were diagnosed with a new episode of major depression, treated with antidepressant medication, and who remained on an antidepressant medication treatment. | NCQA Contact Information: www.ncqa.org | NA | NA | Hedis 2012- Effectiveness of Care: Antidepressant Medication Management: % of members who had a new diagnosis of depression and treated with an antidepressant who remained on antidepressant for acute phase and recovery phase of treatment | NA |
| NQF 0086 PQRI 12 | Title: Primary Open Angle Glaucoma (POAG): Optic Nerve Evaluation Description: Percentage of patients aged 18 years and older with a diagnosis of POAG who have been seen for at least two office visits who have an optic nerve head evaluation during one or more office visits within 12 months. | AMA-PCPI Contact Information: cpe@ama-assn.org | NA | NA | NA | NA |

Note: Each provider must report on 3 core measures (or alternate core) and 3 additional measures.

| NQF Measure Number & PQRI Implementation Number | Clinical Quality Measure Title & Description | Clinical Quality Measure Steward & Contact Information | Core Clinical Quality Measure | UDS Overlap | Health Home Measure | NCQA Diabetes Recognition Program Measures |
|---|--|--|-------------------------------|---|---|---|
| NQF 0088 PQRI 18 | <p>Title: Diabetic Retinopathy: Documentation of Presence or Absence of Macular Edema and Level of Severity of Retinopathy Description: Percentage of patients aged 18 years and older with a diagnosis of diabetic retinopathy who had a dilated macular or fundus exam performed which included documentation of the level of severity of retinopathy and the presence or absence of macular edema during one or more office visits within 12 months.</p> | <p>AMA-PCPI Contact Information: cpe@ama-assn.org</p> | NA | NA | NA | NA |
| NQF 0089 PQRI 19 | <p>Title: Diabetic Retinopathy: Communication with the Physician Managing Ongoing Diabetes Care Description: Percentage of patients aged 18 years and older with a diagnosis of diabetic retinopathy who had a dilated macular or fundus exam performed with documented communication to the physician who manages the ongoing care of the patient with diabetes mellitus regarding the findings of the macular or fundus exam at least once within 12 months.</p> | <p>AMA-PCPI Contact Information: cpe@ama-assn.org</p> | NA | NA | NA | NA |
| NQF 0047 PQRI 53 | <p>Title: Asthma Pharmacologic Therapy Description: Percentage of patients aged 5 through 40 years with a diagnosis of mild, moderate, or severe persistent asthma who were prescribed either the preferred long-term control medication (inhaled corticosteroid) or an acceptable alternative treatment.</p> | <p>AMA-PCPI Contact Information: cpe@ama-assn.org</p> | NA | 2011 UDS Measure. Table 6b, Section H, row 16. | Use of Appropriate Medications for People with Asthma: Hedis 2012-Effectiveness of Care: % pf of members who are identified with persistent asthma and who were appropriately prescribed preferred asthmas medication | NA |
| NQF 0001 PQRI 64 | <p>Title: Asthma Assessment Description: Percentage of patients aged 5 through 40 years with a diagnosis of asthma and who have been seen for at least 2 office visits, who were evaluated during at least one office visit within 12 months for the frequency (numeric) of daytime and nocturnal asthma symptoms.</p> | <p>AMA-PCPI Contact Information: cpe@ama-assn.org</p> | NA | NA | NA | NA |
| NQF 0002 PQRI 66 | <p>Title: Appropriate Testing for Children with Pharyngitis Description: Percentage of children 2-18 years of age who were diagnosed with pharyngitis, dispensed an antibiotic and received a group A streptococcus (strep) test for the episode.</p> | <p>NCQA Contact Information: www.ncqa.org</p> | NA | NA | NA | NA |
| NQF 0387 PQRI 71 | <p>Title: Oncology Breast Cancer: Hormonal Therapy for Stage IC-IIIC Estrogen Receptor/Progesterone Receptor (ER/PR) Positive Breast Cancer Description: Percentage of female patients aged 18 years and older with Stage IC through IIIC, ER or PR positive breast cancer who were prescribed tamoxifen or aromatase inhibitor (AI) during the 12-month reporting period.</p> | <p>AMA-PCPI Contact Information: cpe@ama-assn.org</p> | NA | NA | NA | NA |
| NQF 0385 PQRI 72 | <p>Title: Oncology Colon Cancer: Chemotherapy for Stage III Colon Cancer Patient Description: Percentage of patients aged 18 years and older with Stage IIIA through IIIC colon cancer who are referred for adjuvant chemotherapy, prescribed adjuvant chemotherapy, or have previously received adjuvant chemotherapy within the 12-month reporting period.</p> | <p>AMA-PCPI Contact Information: cpe@ama-assn.org</p> | NA | NA | NA | NA |
| NQF 0389 PQRI 102 | <p>Title: Prostate Cancer: Avoidance of Overuse of Bone Scan for Staging Low Risk Prostate Cancer Patients Description: Percentage of patients, regardless of age, with a diagnosis of prostate cancer at low risk of recurrence receiving interstitial prostate brachytherapy, OR external beam radiotherapy to the prostate, OR radical prostatectomy, OR cryotherapy who did not have a bone scan performed at any time since diagnosis of prostate cancer.</p> | <p>AMA-PCPI Contact Information: cpe@ama-assn.org</p> | NA | NA | NA | NA |
| NQF 0027 PQRI 115 | <p>Title: Smoking and Tobacco Use Cessation, Medical assistance: a. Advising Smokers and Tobacco Users to Quit, b. Discussing Smoking and Tobacco Use Cessation Medications, c. Discussing Smoking and Tobacco Use Cessation Strategies Description: Percentage of patients 18 years of age and older who were current smokers or tobacco users, who were seen by a practitioner during the measurement year and who received advice to quit smoking or tobacco use or whose practitioner recommended or discussed smoking or tobacco use cessation medications, methods or strategies.</p> | <p>NCQA Contact Information: www.ncqa.org</p> | NA | Similar to UDS table 6a, line 26c smoke and tobacco use cessation counseling. | NA | <p>Title: Smoking Status Description: Document smoking status. If smoker, record date that documents smoking cessation counseling or treatment with 12 month abstraction period.</p> |

Note: Each provider must report on 3 core measures (or alternate core) and 3 additional measures.

| NQF Measure Number & PQRI Implementation Number | Clinical Quality Measure Title & Description | Clinical Quality Measure Steward & Contact Information | Core Clinical Quality Measure | UDS Overlap | Health Home Measure | NCQA Diabetes Recognition Program Measures |
|---|---|---|-------------------------------|---|---|---|
| NQF 0055 PQRI 117 | Title: Diabetes: Eye Exam Description: Percentage of patients 18 -75 years of age with diabetes (type 1 or type 2) who had a retinal or dilated eye exam or a negative retinal exam (no evidence of retinopathy) by an eye care professional. | AMA-PCPI Contact Information: cpe@ama-assn.org | NA | NA | NA | Title: Eye Exam Description: Proportion of patients 18 -75 years of age with diabetes (type 1 or type 2) who had a retinal or dilated eye exam or a negative retinal exam (no evidence of retinopathy) by an eye care professional. Record date of most recent retinal or dilated eye exam performed within the 12-month abstraction period. |
| NQF 0062 PQRI 119 | Title: Diabetes: Urine Screening Description: Percentage of patients 18 - 75 years of age with diabetes (type 1 or type 2) who had a nephropathy screening test or evidence of nephropathy. | AMA-PCPI Contact Information: cpe@ama-assn.org | NA | NA | NA | NA |
| NQF 0056 PQRI 163 | Title: Diabetes: Foot Exam Description: The percentage of patients aged 18 – 75 years with diabetes (type 1 or type 2) who had a foot exam (visual inspection, sensory exam with monofilament, or pulse exam). | NCQA Contact Information: www.ncqa.org | NA | NA | NA | Title: Foot Exam Description: The proportion of patients aged 18 – 75 years with diabetes (type 1 or type 2) who had a foot exam. Record date of most recent foot exam within the 12-month abstraction period |
| NQF 0074 PQRI 197 | Title: Coronary Artery Disease (CAD): Drug Therapy for Lowering LDL-Cholesterol Description: Percentage of patients aged 18 years and older with a diagnosis of CAD who were prescribed a lipid-lowering therapy (based on current ACC/AHA guidelines). | AMA-PCPI Contact Information: cpe@ama-assn.org | NA | 2012: New Measure, Table 6B, Section I, Line 17. | NA | NA |
| NQF 0084 PQRI 200 | Title: Heart Failure (HF): Warfarin Therapy Patients with Atrial Fibrillation Description: Percentage of all patients aged 18 years and older with a diagnosis of heart failure and paroxysmal or chronic atrial fibrillation who were prescribed warfarin therapy | AMA-PCPI Contact Information: cpe@ama-assn.org | NA | NA | NA | NA |
| NQF 0073 PQRI 201 | Title: Ischemic Vascular Disease (IVD): Blood Pressure Management Description: Percentage of patients 18 years of age and older who were discharged alive for acute myocardial infarction (AMI), coronary artery bypass graft (CABG) or percutaneous transluminal coronary angioplasty (PTCA) from January 1- November 1 of the year prior to the measurement year, or who had a diagnosis of ischemic vascular disease (IVD) during the measurement year and the year prior to the measurement year and whose recent blood pressure is in control (<140/90 mmHg). | NCQA Contact Information: www.ncqa.org | NA | NA | NA | NA |
| NQF 0068 PQRI 204 | Title: Ischemic Vascular Disease (IVD): Use of Aspirin or Another Antithrombotic Description: Percentage of patients 18 years of age and older who were discharged alive for acute myocardial infarction (AMI), coronary artery bypass graft (CABG) or percutaneous transluminal coronary angioplasty (PTCA) from January 1- November 1 of the year prior to the measurement year, or who had a diagnosis of ischemic vascular disease (IVD) during the measurement year and the year prior to the measurement year and who had documentation of use of aspirin or another antithrombotic during the measurement year. | NCQA Contact Information: www.ncqa.org | NA | 2012: New Measure, Table 6B, Section J, Line 18. | NA | NA |
| NQF 0004 | Title: Initiation and Engagement of Alcohol and Other Drug Dependence Treatment: (a) Initiation, (b) Engagement Description: The percentage of adolescent and adult patients with a new episode of alcohol and other drug (AOD) dependence who initiate treatment through an inpatient AOD admission, outpatient visit, intensive outpatient encounter or partial hospitalization within 14 days of the diagnosis and who initiated treatment and who had two or more additional services with an AOD diagnosis within 30 days of the initiation visit | NCQA Contact Information: www.ncqa.org | NA | NA | Initiation and Engagement of Alcohol and other Drug Dependence Treatment : Hedis 2012-Access/Availability of Care: The % of adolescent and adult patients with a new episode of alcohol and other drug (AOD) dependence who initiate treatment through an inpatient AOD admission, outpatient visit, intensive outpatient encounter or partial hospitalization within 14 days of the diagnosis and who initiated treatment and who had two or more additional services with an AOD diagnosis within 30 days of the initiation visit | NA |
| NQF 0012 | Title: Prenatal Care: Screening for Human Immunodeficiency Virus (HIV) Description: Percentage of patients, regardless of age, who gave birth during a 12-month period who were screened for HIV infection during the first or second prenatal care visit. | AMA-PCPI Contact Information: cpe@ama-assn.org | NA | 2011 UDS Measure. Similar to Table 7, Section A, Line 0, HIV Positive Pregnant Women. | NA | NA |

Note: Each provider must report on 3 core measures (or alternate core) and 3 additional measures.

| NQF Measure Number & PQRI Implementation Number | Clinical Quality Measure Title & Description | Clinical Quality Measure Steward & Contact Information | Core Clinical Quality Measure | UDS Overlap | Health Home Measure | NCQA Diabetes Recognition Program Measures |
|---|---|---|-------------------------------|--|--|---|
| NQF 0014 | Title: Prenatal Care: Anti-D Immune Globulin Description: Percentage of D (Rh) negative, unsensitized patients, regardless of age, who gave birth during a 12-month period who received anti-D immune globulin at 26-30 weeks gestation. | AMA-PCPI Contact Information: cpe@ama-assn.org | NA | NA | NA | NA |
| NQF 0018 | Title: Controlling High Blood Pressure Description: The percentage of patients 18-85 years of age who had a diagnosis of hypertension and whose BP was adequately controlled during the measurement year | NCQA Contact Information: www.ncqa.org | NA | 2011 UDS Measure. Similar to Table 7, Section B, Hypertension. | NA | NA |
| NQF 0032 | Title: Cervical Cancer Screening Description: Percentage of women 21-64 years of age, who received one or more Pap tests to screen for cervical cancer | NCQA Contact Information: www.ncqa.org | NA | 2011 UDS Measure. Similar to Table 6b, Row 11 Pap Tests. | NA | NA |
| NQF 0033 | Title: Chlamydia Screening for Women Description: Percentage of women 15- 24 years of age who were identified as sexually active and who had at least one test for Chlamydia during the measurement year | NCQA Contact Information: www.ncqa.org | NA | NA | Chlamydia Screening for Women: Hedis 2012: Effectiveness of Care: % of women who were identified as sexually active and who had at least one test for Chlamydia | NA |
| NQF 0036 | Title: Use of Appropriate Medications for Asthma Description: Percentage of patients 5 - 50 years of age who were identified as having persistent asthma and were appropriately prescribed medication during the measurement year. Report three age stratifications (5-11 years, 12-50 years, and total). | NCQA Contact Information: www.ncqa.org | NA | NA | Hedis 2012-Effectiveness of Care: % pf of members who are identified with persistent asthma and who were appropriately prescribed preferred asthmas medication | NA |
| NQF 0052 | Title: Low Back Pain: Use of Imaging Studies Description: Percentage of patients with a primary diagnosis of low back pain who did not have an imaging study (plain x-ray, MRI, CT scan) within 28 days of diagnosis | NCQA Contact Information: www.ncqa.org | NA | NA | NA | NA |
| NQF 0075 | Title: Ischemic Vascular Disease (IVD): Complete Lipid Panel and LDL Control Description: Percentage of patients 18 years of age and older who were discharged alive for acute myocardial infarction (AMI), coronary artery bypass graft (CABG) or percutaneous transluminal angioplasty (PTCA) from January 1-November1 of the year prior to the measurement year, or who had a diagnosis of ischemic vascular disease (IVD) during the measurement year and the year prior to the measurement year and who had a complete lipid profile performed during the measurement year and whose LDL-C<100 mg/dL. | NCQA Contact Information: www.ncqa.org | NA | NA | Cholesterol Testing for patient with Cardiovascular Conditions: Hedis 2012: Effectiveness of Care: % of members who were discharged alive for AMI, CABG or PCI or who have a diagnosis of IVD and who had at least one LDL-screening | NA |
| NQF 0575 | Title: Diabetes: Hemoglobin A1c Control (<8.0%) Description: The percentage of patients 18-75 years of age with diabetes (type 1 or type 2) who had hemoglobin A1c <8.0%. | NCQA Contact Information: www.ncqa.org | NA | 2011 UDS Measure. Similar to Table 7, Section C. | Not an exact match to NQF 0575 Hedis 2012 Effectiveness of Care Comprehensive Diabetes Care (HbA1c test and LDL-test) Percentage of members with diabetes who had at least one HbAc test and at least one LDL-C test | Title: HbA1c Testing & Control (>9.0%, <8.0%, <7.0%) Description: The proportion of a patient panel that falls within the stated levels. Date and value of most recent HbA1c performed within the 12-month abstraction period. |

Note: Each provider must report on 3 core measures (or alternate core) and 3 additional measures.